

The Anticoag Management JPP is used by registered nurses in practices and clinics who manage anticoagulation therapy for their patients. This flyer demonstrates how to initiate and use the AMB Anticoag Management Joint Practice Protocol.

AMB Anticoag Management Joint Practice Protocol (JPP)

The protocol can be proposed by the Anticoagulation Clinic MA, RN, Pharmacist, or initiated by the Anticoag Clinic Provider.

➤ Ordering the Anticoag Management JPP




STEP 1: From the Primary Care, Cardiology, or Urology Quick Order Page, the Anticoag Management JPP displays below the AMB Joint Practice Protocols.

STEP 2: Select the **AMB Anticoag Management Joint Practice Protocol** order.

AMB Anticoag Management Joint Practice Protocol

STEP 3: When the MA, RN, or Pharmacist is initiating or renewing the protocol, the **Ordering Physician** window opens.

STEP 4: Select **Proposal**.



STEP 5: Enter the ordering provider's name in the **Physician name** field.

STEP 6: Select Communication type as **Proposal, Cosign Required**.

STEP 7: Select **OK**.


STEP 8: Click the **signature tray**.



STEP 9: The **Orders for Signature** window opens to associate the appropriate diagnosis.

STEP 10: Click **Sign**.

STEP 11: Complete the details within the order.



STEP 12: Click **Sign**.

NOTE: The proposed AMB Anticoag Management Joint Practice Protocol order appears in the ordering provider's message center under Proposed Orders.

The Anticoag Management JPP is active once the ordering provider signs the order.

Renewing Anticoag Management JPP

➤ A reminder populates the **Recommendations** component to renew the **Anticoag Management JPP**.

The screenshot shows a 'Recommendations' window with a '+ ' button in the top right. Below the title bar are two tabs: 'Pending' (selected) and 'Not Due / Historical'. A table below the tabs has columns for 'Recommendation', 'Next Due', 'Recurrence', and 'Orders'. The first row contains 'Anticoag JPP (Anticoag Program Use Only)', 'Today', 'Every 1 YR', and a button labeled 'AMB Anticoag JPP Order'.

AMB JPP Anticoag Management PowerPlan

The **AMB JPP Anticoag Management PowerPlan** is used to order labs and review the **Warfarin Nomogram**.

➤ Placing orders per JPP for **AMB JPP Anticoag Management**.

STEP 1: Select **AMB JPP Anticoag Management PowerPlan**.

STEP 2: Click the **signature tray**.

STEP 3: Associate the appropriate diagnosis in the **Order for Signature** window.

STEP 4: For MA/RNs, in the **Ordering Physician** window:

- Select **Order**.
- Enter the ordering physicians name in the **Physician Name** field.
- Select **Joint Practice Protocol, Active** as the Communication type.
- Select **OK**.

The screenshot shows a 'Joint Practice Protocol' window with a search bar and a list of protocols. The list includes 'AMB Joint Practice Protocol', 'AMB Joint Practice Protocols 17 and younger', 'AMB Joint Practice Protocols 18 and older', 'Anticoag Management JPP', 'Anticoagulation Patient', 'AMB Anticoag Management Joint Practice Protocol This JPP to be utilized', 'AMB JPP Anticoag Management AMB JPP Anticoag Management', and 'Anticoag Management Ortho Joint Practice Protocol'. The 'AMB JPP Anticoag Management' item is highlighted with a red box.

The screenshot shows an 'Ordering Physician' window with a 'P' icon in the top left and a close button in the top right. It has radio buttons for 'Order' (selected) and 'Proposal'. Below are fields for 'AMB JPP Adult', '*Physician name' (with a search icon), and '*Order Date/Time' (with date and time pickers). A '*Communication type' list includes 'Administrative', 'Verbal-Requires Read Back', 'Fax', 'Telephone-Requires Read Back', 'Joint Practice Protocol, Active' (highlighted in blue), 'Patient Care Protocol, Cosign Required', 'Proposal, Cosign Required', and 'Protocol'. 'OK' and 'Cancel' buttons are at the bottom.

From the Office of Clinical Informatics
AMB Anticoag Management Joint Practice Protocol
January 16, 2024
Page 3 of 3

STEP 6: The AMB JPP Anticoag Management PowerPlan opens.

STEP 7: Select the appropriate order(s) by clicking the checkbox and complete any missing details.

Laboratory:
Patient on anticoagulation therapy presents for testing. Use PT (Prottime) INR lab order for testing outside the office. Use INR POC 85610 order for point of care testing

Order	Frequency
<input checked="" type="checkbox"/> INR POC 85610	
<input type="checkbox"/> PT (Prottime) INR	BLOOD, Stat, ONCE
<input type="checkbox"/> CBC may be ordered at baseline, annually, for clinical suspicion of anemia or occult bleeding, or for new onset or worse than baseline shortness of breath, pallor, or fatigue	
<input type="checkbox"/> HGB less than 7 g/dL: Refer patient to Emergency Department immediately	
<input type="checkbox"/> CBC without Differential	BLOOD, Routine, ONCE
<input type="checkbox"/> The following may be ordered at baseline, annually, and as directed by DOAC Management Workflow	
<input type="checkbox"/> Comprehensive Metabolic Panel (CMP)	BLOOD, Routine, ONCE
<input type="checkbox"/> Creatinine	BLOOD, Routine, ONCE
<input type="checkbox"/> Hepatic Function Panel	BLOOD, Routine, ONCE

Medication:
Medication Refill: Anticoagulant medication managed by the clinic through a referral or managed by primary care may be refilled by AC nurse, AC pharmacist or primary care nurse up to 6 months per refill. The anticoagulant medication must be present on the current medication list in the patient's medical record and the patient must maintain appropriate follow-up

Order	Frequency
<input type="checkbox"/> Administer to those patients requesting the flu vaccination, if not previously administered.	
<input type="checkbox"/> influenza virus vaccine, inactivated (Fluarix PF Quadri...)	0.5 mL, Susp, IM, Vaccine, 1 Doses/Times
<input type="checkbox"/> influenza virus vaccine, inactivated (Fluzone High-Do...)	Select an order sentence

Non Categorized
Providers must exercise independent clinical judgment when using order sets.

Details
Dx Table Orders For Co-signature Save as My Favorite Plan for Later **Initiate Now** Cancel

STEP 8: Select **Initiate Now**.

STEP 9: Click **Orders for Signature**.

STEP 10: Click **Sign**.

NOTE: The AMB Joint Practice Protocol Order must be signed and active before an MA or RN can place orders per JPP.

DO NOT choose orders from the POC component. POC component orders do not have the reference text criteria built into them to be ordered per JPP.